

An Overview of the Bioactive Constituents, Biological Functions, and Pharmacological Impacts of *Viscum album* L.

Eva Kleszken¹, Adrian Vasile Timar², Adriana Ramona Memete¹, Florina Miere (Groza)³,
Simona Ioana Vicas^{2*}

¹Doctoral School of Biomedical Science, University of Oradea, Oradea, Romania.

²Department of Food Engineering, University of Oradea, Oradea, Romania.

³Faculty of Medicine and Pharmacy, University of Oradea, Oradea, Romania.

*E-mail ✉ svicas@uoradea.ro

Received: 03 August 2022; Revised: 24 October 2022; Accepted: 26 October 2022

ABSTRACT

The *Viscum* genus includes a variety of species found in regions such as Europe, Africa, Asia, the Americas, and Australia. Mistletoe extracts and their formulations are widely used in alternative and complementary medicine for treating a wide range of ailments. This review focuses on studies that examine the phytochemical properties of mistletoe, which vary depending on the host tree and highlights research on its therapeutic effects. Mistletoe contains an array of bioactive compounds, including lectins, viscotoxins, phenolic acids, flavonoids, alkaloids, terpenoids, and polysaccharides. These compounds contribute to mistletoe's diverse biological activities, such as anticancer, anti-inflammatory, and cardiovascular benefits, while also helping to alleviate chemotherapy side effects and enhance immune response. This review emphasizes the relationship between the host tree and the bioactive components of mistletoe, especially the lectins, viscotoxins, and phenolic compounds such as flavonoids and phenolic acids. The therapeutic properties of mistletoe are further investigated by examining the subspecies and host trees, and various patents based on mistletoe applications are also discussed. Mistletoe is introduced as a valuable medicinal plant, with high potential for further investigation in the development of targeted therapies.

Keywords: Anticancer, *Viscum album*, Flavonoids, Host trees, Phenolic acid, Antioxidant

How to Cite This Article: Kleszken E, Timar AV, Memete AR, Miere (Groza) F, Vicas SI. An Overview of the Bioactive Constituents, Biological Functions, and Pharmacological Impacts of *Viscum album* L. Spec J Pharmacogn Phytochem Biotechnol. 2022;2:53-72. <https://doi.org/10.51847/pPFuHQUG2R>

Introduction

Parasitic angiosperms comprise over 4,700 species distributed across 277 genera, with mistletoe representing 87 genera and at least 1,670 species. Mistletoe is classified as a hemiparasite, relying on various tree and shrub hosts, and resembling an extensively developed shrub [1, 2]. Parasitic interactions, whether as a parasite or a host, are vital for maintaining ecosystem balance and promoting biological evolution [3, 4].

Mistletoe (*Viscum album* L.), an evergreen plant, depends on its host for water and nutrients but also produces its carbohydrates through photosynthesis. It is commonly found on both coniferous and deciduous trees [2]. This large hemiparasitic species is found in regions of Europe, Africa, the Americas, Australia, and Asia, exhibiting diverse habits, preferred hosts, and morphologies based on geographical location [5-7].

Pharmacological studies have identified numerous bioactive compounds in mistletoe, including lectins, viscotoxins, lignans, flavonoids, amines, and polysaccharides [8, 9]. Among these, flavonoids and phenolic acids act as potent antioxidants and are involved in the plant's biological effects, particularly in the prevention of diseases like cancer caused by oxidative stress and free radicals in the body [10-13]. Mistletoe has demonstrated therapeutic value in treating various types of cancer, including those affecting the breast [14, 15], pancreas [16], larynx [17], bladder [18], and leukemia [19]. Over time, additional bioactive effects of mistletoe have been

observed, including its impact on neurological disorders, its antiviral and antibacterial properties, anti-inflammatory actions, antiepileptic effects, and immune system stimulation [2, 11, 13, 20].

Due to its bioactive compounds and diverse health benefits, mistletoe has become the subject of numerous studies, leading to the creation of several pharmaceutical preparations, such as Iscador, Isorel, Iscucin, Lektinol, Eurixor, Helixor, Abnoba-Viscum, and recombinant lectin ML-1 [13, 21].

This review focuses on the relationship between the host tree and the bioactive constituents of mistletoe, particularly lectins, viscotoxins, flavonoids, and phenolic acids. It also outlines the potential therapeutic effects of mistletoe according to its subspecies and host tree and discusses the various patents related to mistletoe-based applications.

Materials and Methods

The review follows the PRISMA 2020 flowchart methodology recommended by Page *et al.* [22], and the steps, along with selection criteria, are illustrated in **Figure 1**. The literature was gathered from PubMed, Scopus, Science Direct, Elsevier, Google Scholar, and Google Patents. Search terms included “*Viscum album*”, “*V. coloratum*”, “*V. articulatum*”, “mistletoe viscotoxin”, “mistletoe lectin”, “mistletoe effects”, “mistletoe patents”, “mistletoe compounds”, and “mistletoe history”. Only studies in English and Romanian were considered. A total of 100 relevant studies were selected and included in this review.




Figure 1. PRISMA 2020 flow diagram for the present review

Results and Discussion

The historical context of mistletoe

Over 150 years ago, Charles Darwin took note of the co-evolutionary dynamics between mistletoe and the birds that interact with it, sparking significant scientific interest. In addition, the interactions between herbivorous

insects and plants, which can facilitate transitions between plant species, also gained the attention of researchers. Mistletoe is considered an ancestral host for certain butterfly species [23-25].

The number of host species affected by mistletoe plays a crucial role in determining the parasite's distribution, prevalence, and virulence. However, macroecological studies of this relationship across various regions remain scarce. Over time, human activities and environmental changes, including hybridization, have influenced the dynamics of parasitic species [1, 26, 27].

Mistletoe's remarkable therapeutic effects were later discovered, and it gradually became integrated into traditional medicine, especially species like *V. album* and *Loranthus* (*Taxillus Chinensis* (DC.) Danser) [28, 29]. The latter species, native primarily to southern and southwestern China, is known as "Sang Ji Sheng" in traditional Chinese medicine, where its stems and leaves are used to treat conditions like rheumatism, arrhythmia, hypertension, stroke, and angina pectoris [28]. *V. album*, with its longstanding use in German-speaking regions for cancer treatment, has a similar historical significance, spanning nearly a century [29].

Beyond its medicinal properties, mistletoe is also surrounded by numerous legends and beliefs that have been passed down through generations and continue to be remembered today. The plant has been thought to bring good fortune, promote healing, and attract angels. Its mystical reputation dates back to ancient times. The Celts, for instance, believed that mistletoe, created by a bolt of lightning striking a tree, possessed magical qualities. They highly valued mistletoe growing on oak trees, referring to it as the "Oak Tear," and the act of harvesting this plant was a ceremonial event. Only after the winter solstice would the mistletoe be carefully cut with a golden sickle by a priest dressed in white. Mistletoe was also seen as a symbol of femininity (*Naturalis Historia*, books XVI, XXIV, XXXIII).

The Greeks regarded mistletoe as a symbol of the underworld, associating it with Aeneas' key to the underworld. For the Romans, it was a fertility symbol, often worn as a crown by the goddess Diana. In Scandinavian traditions, mistletoe symbolizes peace, love, harmony, and spiritual purification. It was also believed to ward off evil spirits. In European folklore, mistletoe holds a special place. Around Christmas and New Year's, it was common for people to hang mistletoe branches on doorways and windows to attract good health and fortune. Another legend involves enemies reconciling under the mistletoe, a custom later adopted by lovers, who would exchange a kiss beneath it as a pledge of love, symbolizing happiness and fulfillment. The town of Tenbury Wells in the UK hosts an annual mistletoe festival, where a Mistletoe Queen is crowned, wearing a mistletoe crown [30].

Botanical description

V. album is a hemiparasitic shrub that grows on the branches of trees. It develops slowly, with its growth increasing yearly, marking its age. This plant can live up to 70 years [31, 32], often growing into a large bush up to one meter in diameter (**Figure 2a**). The mistletoe attaches firmly to the host tree, sending a well-developed axis into the tree's bark, penetrating the woody tissue. This structure extends into the host tree, forming a root-like system (haustorium) that absorbs water, sugars, amino acids, and minerals, while also synthesizing its own primary and secondary metabolites [33].

The stem of mistletoe is cylindrical, thick, and dichotomously branched, ranging from 30-60 cm in width, growing thicker near the nodes, and can be easily torn. It is yellow-green. Its leaves are evergreen, arranged in opposite pairs, without stalks, and possess entire edges. The leaves have a ribbon-like appearance, thick and narrow, and display a yellow to intense green color. They also show 4-5 veins on the underside (**Figure 2b**). The size of the leaves varies based on the host tree and harvest time. Leaves gathered in April, especially those from mistletoe on acacia trees, are larger. These leaves emit a specific scent and have a bitter-sour taste.

The optimal harvesting period for mistletoe is from November to April. Its flowers are small (2-3 mm in diameter), yellow-greenish, and dioecious, blooming in March-April (**Figure 2c**). The fruits of mistletoe are spherical, with a translucent, glassy appearance, approximately 8 mm in diameter. They are green when young and turn white and translucent from September to October (**Figure 2d**), but are toxic from November to March (**Figure 2f**). Typically, the fruits, containing 1-2 seeds surrounded by a sticky, jelly-like substance, are harvested in clusters between September and December (**Figure 2e, 2g**).

Botanists mistakenly identified *V. album* as oak mistletoe (*L. europaeus*) up until the 18th century. *V. album* belongs to the Viscaceae family, while *L. europaeus* belongs to the Loranthaceae family. Both species are part of the Santalales order [7, 34].

L. europaeus has young twigs that are brownish, lance-shaped leaves, and short, entire edges. It sheds its leaves in winter, unlike *V. album*, which retains its green leaves throughout the year. *L. europaeus* blooms between May

and June, with small yellow-greenish leaves, whereas *V. album* produces small clusters of yellowish fruits, ripening in October, which are dispersed by birds [35].




Figure 2. Morphology of *V. album*. a) Mistletoe (*V. album* on acacia), b) leaves of mistletoe, c) Flower of mistletoe, d) Unripe fruits of mistletoe, e) Unripe fruits with seeds, f) White fruits of mistletoe, g) White fruits with seeds

Taxonomic classification

The Viscaceae family includes seven genera: *Viscum*, *Phoradendron*, *Notothixos*, *Korthalsella*, *Ginalloa*, *Dendrophthora*, and *Arceuthobium*. Traditionally, *V. album*, commonly known as mistletoe, was classified under the *Viscum* genus within the Viscaceae family. However, recent genetic studies and the updated APG III (Angiosperm phylogeny group) system suggest that mistletoe is more appropriately placed within the Santalaceae family. The *Viscum* genus consists of approximately 90 species, with about two-thirds found in Africa and one-third in both Eurasia and Africa [36].

The *Phoradendron* genus, which includes the North American oak mistletoe (*Phoradendron serotinum*), is primarily found in North and South America, extending to Argentina. *Loranthus europaeus*, a species of mistletoe from the Loranthaceae family, is widespread in the Americas. The *Notothixos* genus is native to eastern Australia and its surrounding islands. *Korthalsella* species are primarily found in East Asia, Malaysia, Australia, and New Zealand. The *Ginalloa* genus contains a small number of species, located mostly in Southeast Asia. The *Dendrophthora* genus, like *Phoradendron*, is found in Central America. The *Arceuthobium* genus consists of around 24 species, distributed across the USA and Mexico [37].

Host trees and geographical distribution

The European mistletoe is classified into various subspecies, which can be distinguished based on their host trees, as these subspecies are often morphologically quite similar. Differences in leaf shape and size, fruit color, and the specific host tree help in identifying these mistletoe species. *V. album* parasitizes over 400 host species [38]. **Table 1** provides a summary of existing literature on mistletoe species and subspecies, detailing their host trees across different continents, leaf size, and fruit color.

Table 1. Mistletoe species from different continents, their host trees, leaf size, and fruit color

Mistletoe species	Spread	Host tree	Leaves	Fruits	References
Europe					
<i>V. album</i> ssp. <i>abietis</i>	Central Europe	Fir (<i>Abies alba</i>)	Up to 8 cm	Whites	[7]
<i>V. album</i> ssp. <i>album</i>	Europe, Southwest Asia	Apple(<i>Malus</i>), Linden (<i>Tilia</i>), Willow (<i>Salix vitellina</i>), Poplar (<i>Populus nigra</i>), Oak (<i>Quercus</i>) Hawthorn (<i>Crataegus monogyna</i>), Apricot (<i>Prunus armeniaca</i>), Acacia (<i>Robinia pseudoacacia</i>) It never appears on Beech.	Up to 5 cm	Whites	[7]
<i>V. album</i> ssp. <i>austriacum</i>	Central Europe, rarely in Greece	Coniferous (<i>Larix</i> , <i>Pinus</i> , <i>Picea</i>).	4-6 cm	Yellow	[7]
<i>V. album</i> ssp. <i>Creticum</i>	Eastern Crete	Pine (<i>Pinus</i>)	Small	Whites	[5]
Asia					
<i>V. album</i> ssp. <i>meridianum</i>	Southeast Asia	Hornbeam (<i>Carpinus monbeigiana</i>), Ash (<i>Fraxinus</i>), Maple (<i>Acer</i>), Walnut(<i>Juglans regia</i>), Plum(<i>Prunus pseudocirasus</i>) Mountain ash (<i>Sorbus megalopa</i>)	3-5 cm	Yellow	[39]
<i>V. album</i> ssp. <i>coloratum</i>	China	Oak (<i>Quercus</i> sp.).	2-4 cm	Red	[40-42]
Africa					
The red mistletoe (<i>V. cruciate</i>).	North Africa (Morocco, Libya) ends in South Africa, Rarely in South-western Spain, southern Portugal	Olive (<i>Olea europaea</i>)	Small	Red	[43]
<i>L. ferrugineus</i>	Africa	Oak (<i>Quercus</i> sp.), Acacia (<i>Robinia pseudoacacia</i>) Euphorbia	4-8 cm	Yellow	[44]
<i>L. micranthus</i>	Nigeria	Oak (<i>Quercus</i>),	4-6 cm	Yellow	[45]
Australia					
<i>Amyema maidenii</i> ssp. <i>maidenii</i>	Australia	<i>Eucalyptus</i>	Flat	Yellow	[46]
<i>A. gibberula</i> ssp. <i>gibberula</i>			Long and cylindrical	Whites	[46]
<i>A. bifurcata</i> ssp. <i>bifurcata</i>			Flat	Red	[47]
<i>Lysiana exocarpi</i> ssp. <i>exocarpi</i>		Acacia	3-15 cm	Red or black	[48]
<i>L. murrayi</i>			2.5–6 cm	Pink or red	[46]
<i>L. spatulata</i>			3-7 cm	Red	[47]

<i>L. subfalcata</i>			2-12 cm	Yellowish	[46]
America					
The American Mistletoe (<i>Phoradendron californicum</i>) and the dwarf mistletoe (<i>Arceuthobium minutissimum</i>)	North, America, California,	Oak (<i>Quercus</i> sp.),	4-8 cm	Red	[49]
	South America, Argentina	Pine (<i>Pinus</i>)	1-2 cm	Whites	

Mistletoe thrives in diverse environments, including woodlands, hillsides, and mountainous regions, where it attaches to various host trees. It is commonly found on deciduous species such as birch, ash, maple, poplar, lime, and willow, as well as on coniferous trees like fir and pine. Additionally, it parasitizes fruit-bearing trees, including apples, pears, plums, and cherries. Species within the *Viscum* genus are distributed across temperate regions of Europe and Asia, as well as tropical and subtropical zones in Africa, Madagascar, Australia, and North and South America.

Bioactive compounds in mistletoe

V. album L. has long been the subject of scientific inquiry due to the bioactive nature of its constituents. Numerous metabolites have been extracted from European mistletoe, some of which are not synthesized within the plant itself but instead originate from the host tree, such as specific alkaloids [50].

A study conducted by Jäger *et al.* [33] identified key biomarkers—including arginine, pipecolic acid, lysine, dimethoxycoumarin, and sinapyl alcohol—associated with *V. album* ssp. *album* growing on three different host species: *Malus domestica*, *Quercus robur*, and *Ulmus carpiniifolia* [33].

Similarly, Zhang *et al.* [51] analyzed the metabolic profile of *V. coloratum* collected from three distinct host species—*Ulmus pumila* L., *Salix babylonica*, and *Populus ussuriensis* Kom.—in two different climatic regions of China (temperate continental and warm temperate humid monsoon climates). Their findings revealed that the synthesis and accumulation of three primary flavanone-class metabolites varied depending on both the host tree and environmental conditions [51].

Polyphenols: structure and classification

Flavonoids, a subgroup of polyphenolic compounds, are secondary metabolites of plant origin with antioxidant properties. In mistletoe leaves, these compounds occur in either free form or as esters and glycosides. Across the plant kingdom, over 4,000 flavonoid variants have been identified [52].

Polyphenols play a crucial role in plant metabolic processes, with approximately 8,000 plant-derived polyphenolic compounds documented for their antioxidant and antitumor effects [53-58]. These organic, water-soluble compounds possess at least one aromatic ring with one or more hydroxyl groups, which contributes to their strong antioxidative capacity [52, 59-61].

Among the polyphenols found in mistletoe, phenolic acids are particularly significant, including hydroxybenzoic acids (such as gallic acid and protocatechuic acid) and hydroxycinnamic acids (such as caffeic acid, ferulic acid, and synaptic acid). Additionally, mistletoe contains various flavonoids, including flavanones (naringenin, eriodictiol), flavones (apigenin), and flavonols (3-O-methylquercetin, myricetin, kaempferol). **Figure 3** provides an overview of the key polyphenols identified in different *V. album* subspecies and their respective host trees.




Figure 3. The classification and chemical structure of the main polyphenols identified in *V. album*

The concentration and profile of flavonoids in mistletoe can differ substantially based on factors such as the host tree, the stage of vegetation, and the timing of harvest [33, 41, 51, 62, 63]. **Table 2** presents the most significant flavonoids identified to date in mistletoe leaves and stems.

Table 2. Polyphenols identified in different subspecies mistletoe leaves or stems depending on the host tree

Species	Host trees	The part used	Flavonoids	Phenolic acids	References
<i>V. album</i> ssp. <i>album</i>	Maple (<i>Acer platanoides</i> L.)	Leaves	Flavone: Apigenin Flavonol: 3-O-Metil Quercetin Flavanones: Naringenin	Hydroxibenzoic acid: Gallic acid, Gentisic acid, Protocatechuic acid, p-hydrobenzoic, Salicylic acid, Cinnamic acid: Caffeic acid, Ferulic acid, p-cumaric acid, Rosmarinic acid, Sinapic acid	[64]
				Hydroxycinnamic acid: Rosmarinic acid	[65]
<i>V. album</i> ssp. <i>album</i>	Jugastru (<i>Acer campestre</i>)	Leaves	Flavonol: Kampherol Quercetin	Hydroxibenzoic acid: Salicylic acid, p-Hydrobenzoic acid, Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid, Ferulic acid, Sinapic acid, trans-Cinnamic acid.	[64]

					Hydroxibenzoic acid: p-Hydrobenzoic acid Salicylic acid, Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid. Ferulic acid, Sinapic acid, trans-Cinnamic acid,	[65]
					Hydroxycinnamic acid: Rosmarinic acid, trans-Cinnamic acid	[65]
					Flavanol: Myricetin Flavonol: 3-O-Metil Quercetin Rhamnazin	[12]
					Hydroxibenzoic acid: Gallic acid, Gentisic acid, p-Hydrobenzoic acid, Salicylic acid, Vanillic acid Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid, Rosmarinic acid,	[64]
<i>V. album</i> ssp. <i>album</i>	Silver coat (<i>Acer saccharinum</i> L.)	Leaves			Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid Syringic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid p-Coumaric acid, Sinapic acid,	[12]
					Flavanones: Eriodictyol Flavonol: 3-O-Metil Quercetin	
<i>V. album</i> ssp. <i>album</i>	Hawthorn (<i>Crataegus monogyna</i>)	Leaves			Hydroxibenzoic acid: Protocatechuic acid, p-Hydrobenzoic acid, Salicylic acid, Syringic acid Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid Ferulic acid, p-Coumaric acid Sinapic acid, trans-Cinnamic acid,	[65]
					Flavonol: Kampherol Quercetin 3-O-Metil Quercetin Flavanones: Eriodictyol	
					Hydroxibenzoic acid: Syringic acid Protocatechuic acid Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid, Ferulic acid, Rosmarinic acid, Sinapic acid, trans-Cinnamic acid,	[65]
					Flavonol: Kampherol	
<i>V. album</i> ssp. <i>album</i>	Ash (<i>Fraxinus excelsior</i> L.)	Stems			Hydroxibenzoic acid: Syringic acid Protocatechuic acid Hydroxycinnamic acid: Caffeic acid, Chlorogenic acid, Ferulic acid, Rosmarinic acid, Sinapic acid, trans-Cinnamic acid,	[65]
					Flavonol: Kampherol	

<i>V. album</i> ssp. <i>album</i>	Ash (<i>Fraxinus pensylvanica</i> Marsh)	Leaves	Flavanones: Eriodictyol Naringenin Flavonol: 3-O-Metil Quercetin Sakuratin Rhamnazin	Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid, Syringic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, Sinapic acid	[64]
			Flavanones: Eriodictyol Naringenin Flavonol: Quercetin 3-O-Metil Quercetin Sakuratin Rhamnazin		[12]
			Flavonol: Quercetin 3-O-Metil Quercetin Myricetin, Sakuratin Isorhamnetin Rhamnetin Rhamnazin Flavanones: Naringenin Eriodictyol		[64]
<i>V. album</i> ssp. <i>album</i>	Apple (<i>Malus Domestica</i> Borkh)	Leaves	Flavonol: Quercetin		[65]
			Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Sakuratin Rhamnazin Flavanones: Naringenin		[12]
				Hydroxibenzoic acid: Gallic acid Vanillic acid, Gentisic acid, Protocatechuic acid, Veratric acid Hydroxycinnamic acid: p-Coumaric acid	[64]
				Hydroxibenzoic acid: p-hydroxybenzoic acid, Protocatechuic acid, Salicylic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, Sinapic acid, Rosmarinic acid,	[65]
				Hydroxibenzoic acid: Syringic acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, Rosmarinic acid Sinapic acid,	[65]
<i>V. album</i> ssp. <i>album</i>	Apple (<i>Malus Domestica</i> Borkh)	Stems			

<i>V. album</i> ssp. <i>album</i>	Poplar (<i>Populus nigra</i> L.)	Leaves	Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Sakuratin, Rhamnazin Flavanones: Naringenin Flavone: Apigenin	[65]
			Hydroxibenzoic acid: Gallic acid, Gentisic acid, Protocatechuic acid, Hydroxycinnamic acid: p-Coumaric acid	[64]
			Hydroxibenzoic acid: Salicylic acid, Protocatechuic acid, p-hydrobenzoic acid, Hydroxycinnamic acid: Ferulic acid, Rosmarinic acid, Sinapic acid,	[65]
			Flavanones: Naringenin Eriodictyol Sakuranetin Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Rhamnazin	[12]
			Hydroxibenzoic acid: p-hydrobenzoic acid, Salicylic acid, Syringic acid Vanillic acid Hydroxycinnamic acid: Caffeic acid, Ferulic acid,	[65]
<i>V. album</i> ssp. <i>album</i>	Poplar (<i>Populus nigra</i> L.)	Stems	Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid Hydroxycinnamic acid: Caffeic acid, Ferulic acid,	[65]
<i>V. album</i> ssp. <i>album</i>	Acacia (<i>Robinia pseudoacacia</i>)	Leaves	Flavonol: Quercetin Kaempferol	[65]
		Stems	Flavonol: Kampherol Hydroxibenzoic acid: Protocatechuic acid,	[65]
<i>V. album</i> ssp. <i>album</i>	Mountain ash (<i>Sorbus aucuparia</i> L.)	Leaves	Flavonol: Quercetin 3-O-Metil Quercetin Isorhamnetin Myricetin, Sakuratin Rhamnazin Flavanones: Naringenin	[12]

			Hydroxibenzoic acid: Gallic acid, Gentisic acid Protocatechuic acid, p –hydroxybenzoic acid Salicylic acid, Syringic acid Vanillic acid, [64] Veratric acid, Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid Sinapic acid,
			Flavanones: Naringenin Eriodictyol Flavonol: Quercetin 3-O-Metil Quercetin [12] Isorhamnetin Sakuratin Rhamnazin
<i>V. album</i> ssp. <i>album</i>	Linden (<i>Tilia cordata</i> Mill)	Leaves	Flavanones: Naringenin Eriodictyol Flavonol: 3-O-Metil Quercetin Isorhamnetin Rhamnazin Flavones: Luteolin Hydroxibenzoic acid: Gallic acid, Protocatechuic acid, p-hydrobenzoic acid, Salicylic acid Syringic acid, Vanillic acid [12] Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid.
<i>V. album</i> ssp. <i>abietis</i>	Fir (<i>Abietis alba</i> Mill)	Leaves	Flavanones: Naringenin Flavonol: 3-O-Metil Quercetin Rhamnazin Rhamnetin Flavone: Apigenin Hydroxibenzoic acid: Salicylic acid, Protocatechuic acid, 4-hydrobenzoic acid, Vanillic acid, [66] Hydroxycinnamic acid: Caffeic acid, Ferulic acid, p-Coumaric acid, Sinapic acid.
<i>V. album</i> ssp. <i>austriacum</i>	Pine (<i>Pinus sylvestris</i> L.)	Leaves	Flavanones: Naringenin Eriodictyol Sakuranetin Flavonol: Quercetin 3-O-Metil Quercetin [12] Isorhamnetin Myricetin, Rhamnazin Rhamnetin, Kampherol
		Leaves fruits seeds	Hydroxibenzoic acid: Protocatechuic acid, Salicylic acid, 4-hydroxybenzoic acid, Vanillic acid, [67] Hydroxycinnamic acid: Caffeic acid, Ferulic acid,

p-Coumaric acid, Sinapic acid				
<i>V. album</i> ssp. <i>austriacum</i>	Guava (<i>Psidium guajava</i>)	Leaves	Flavanones: Naringenin	[68]
	Oak (<i>Quercus crispula</i>)	Leaves stems	Flavanones: Eriodictyol	[19]
<i>V. album</i> ssp. <i>coloratum</i>	Elm (<i>Ulmus pumila</i> L.)			
	Willow (<i>Salix babylonica</i> L.) Poplar (<i>Populus ussuriensis</i> Kom)	Leaves	Flavanones: Eriodictyol	[51]

V. album is a medicinal plant known for its diverse range of phytochemicals. Among its bioactive compounds, quercetin stands out due to its strong antioxidant properties and significant antitumor activity. This flavonol has been identified in multiple subspecies, including *V. album* ssp. *album*, *V. album* ssp. *abietis*, *V. album* ssp. *austriacum*, and *V. album* ssp. *coloratum* [12, 63]. The flavanones Naringenin and Eriodictyol are present in both *V. album* ssp. *album* and *V. album* ssp. *austriacum* [12, 51, 65]. Additionally, caffeic acid, a hydroxycinnamic acid, has been found in various *V. album* subspecies [12, 67].

Lectins and viscotoxins

V. album contains a range of bioactive molecules, from small compounds like phenolic acids and flavonoids to larger proteins such as lectins and viscotoxins [69].

Lectins are proteins or glycoproteins that can selectively reversibly bind to carbohydrates without modifying their structure. They are classified based on their sugar-binding specificity—monospecific lectins interact with a single sugar (e.g., glucose or galactose), while polyspecific lectins can bind multiple sugars [24, 70]. Three distinct lectins, designated ML-1, ML-2, and ML-3 (ML for Mistlelectin), are found in mistletoe, differing in molecular weight and carbohydrate affinity [71-73].

ML-1 consists of two polypeptide chains connected by a disulfide bridge [74]. The A chain comprises three domains with molecular weights of 29 kDa, 27 kDa, and 25 kDa, respectively, and exhibits RNA N-glycosidase activity [75]. The B chain, made up of two domains with molecular weights of 32 kDa and 25 kDa, specifically binds to D-galactose [24, 76, 77]. The B chain's carbohydrate-binding ability is associated with lectins' cytotoxic action, particularly their role in tumor cell targeting. When lectins interact with receptors on cancer cell surfaces, they induce apoptosis, disrupt the cell cycle, and inhibit angiogenesis, leading to tumor suppression [78]. Both chains of ML-1 contribute to its cytotoxic properties [24]. ML-2 binds to D-galactose/N-acetyl-D-galactosamine, while ML-3 is specific to N-acetyl-D-galactosamine [72, 79]. Host tree species influence the composition of lectins in mistletoe, with ML-1 being prevalent in *V. album* growing on oak (*Quercus* spp.) and poplar (*Populus nigra*), while ML-3 is dominant in mistletoe found on conifers like fir (*Abies alba*) and pine (*Pinus sylvestris*) [71].

Viscotoxins are polypeptides with a high cysteine content, consisting of approximately 46 amino acids and three to four disulfide bonds, with molecular weights around 5 kDa [80]. Six structurally distinct viscotoxin isomers—A1, A2, A3, B, B2, C1, and 1-PS—have been identified in *V. album* [72]. These peptides exhibit cytotoxic effects on tumor cells and also play a role in immune modulation [15]. The concentration and composition of lectins and viscotoxins in mistletoe are influenced by the host tree species.

In *V. album* ssp. *album*, viscotoxins A2 and A3 are the predominant types, with 1-PS absent. *V. album* ssp. *austriacum*, on the other hand, contains mostly viscotoxin 1-PS, while A2 and A3 appear in smaller amounts. *V. album* ssp. *abietis* is characterized by the presence of all viscotoxins, with A3 being the most abundant and A2 not detected [81].

Biological activities and medicinal potential

Mistletoe has been shown to exhibit a wide range of biological activities, including anticancer effects, apoptosis induction, antiviral and antibacterial properties, and immune system modulation [2, 11, 20, 82, 83].

Both in vitro and in vivo studies have demonstrated the pharmacological effects of *V. album*. **Table 3** provides an overview of the medicinal activities associated with various mistletoe subspecies (*V. album* ssp., *Viscum coloratum*, and *Viscum articulatum*) about their respective host trees.

Table 3. The pharmacological activity of *V. album* ssp., *Viscum coloratum*, and *Viscum articulatum*

Species/Host tree	Biological activity	Bioactive compounds	Sample type	Type of experiment	References
<i>Viscum album</i> / Apple (<i>Malus domestica</i>)	Anticancer	Lectins	Aqueous Extract + triterpene extract	The human osteosarcoma cell lines 143B and Saos-2	[84]
<i>V. album</i> ssp. <i>coloratum</i> / Poplar (<i>Populus nigra</i>)		Lectins	Aqueous extract	<i>In vitro</i> and <i>vivo</i> -on the growth of melanoma cells in mice.	[85]
<i>V. album</i> ssp. <i>coloratum</i> /ns*		Viscotoxins	Aqueous extract	Randomized clinical trial in patients with locally advanced or metastatic pancreatic cancer	[86]
<i>Viscum album</i> / Apple (<i>Malus domestica</i>), Oak (<i>Quercus</i> spp) Poplar (<i>Populus nigra</i>) Acacia (<i>Robinia pseudo-acacia</i>)				Patient with differentiated squamous cell carcinoma	[87]
<i>V. album</i> L. / Citrus	Antihypertension	Flavonoids	Aqueous extract	Male rats-salt-induced hypertension	[88]
<i>V. album</i> L. /ns*			Ethanollic extract	Wistar rats of both sexes	[89]
<i>V. album</i> /ns*			Ethanollic extract	Hypertensive patient	[90]
<i>V. articulatum</i> / <i>Cordia macledoi</i>		Acid oleanolic	Methanollic extract	Male Wistar rats- dexamethasone-induced hypertension	[91]
<i>V. album</i> / <i>Kola acuminata</i> tree	Antidiabetic		Aqueous extract	Alloxanized male Wistar rats	[92]
			Aqueous extract	STZ-diabetic male Wistar rats	[93]
<i>V. album</i>	Antivirals		Aqueous extract	Human parainfluenza virus type 2 (HPIV-2) growth in Vero cells	[83]
<i>V. album</i> ssp. <i>abietis</i>	Antibacterial		n-hexane extract	<i>In vitro</i> (<i>Bacillus subtilis</i> , <i>Staphylococcus aureus</i> , <i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Enterobacter cloacae</i> , and <i>Proteus vulgaris</i>)	[94]
<i>V. album</i> /ns*			Extracts (1% HCl, ethanol, acetone, and 5% acetic acid)	<i>In vitro</i> (<i>Escherichia coli</i> , <i>Pseudomonas aeruginosa</i> , <i>Staphylococcus aureus</i> , and <i>Klebsiella aerogenes</i>)	[95]
<i>V. album</i> / cocoa and cola trees			Methanollic extract	<i>In vitro</i> (<i>Aspergillus niger</i> , <i>Fusarium exosporium</i> , <i>Penicillium oxalium</i> , and <i>Microsporum canis</i>)	[96]
<i>V. album</i> L. ssp. <i>album</i> /Armeniaca vulgaris Lam. (apricot); <i>V. album</i> ssp. <i>abietis</i> / <i>Abies bornmülleriana</i> Mattf. (fir); <i>V. album</i> ssp. <i>austriacum</i> / <i>Pinus nigra</i> (pine)			Ethanollic extract	<i>In vitro</i> (<i>Mycobacterium tuberculosis</i> H37Ra)	[97]

Antioxidant	Flavonoids	<i>V. album</i> L. ssp. <i>album</i> / <i>Armeniaca vulgaris</i> Lam. (<i>apricot</i>); <i>V. album</i> ssp. <i>abietis</i> / <i>Abies bornmülleriana</i> Mattf. (<i>fir</i>), ; <i>V. album</i> ssp. <i>austriacum</i> / <i>Pinus nigra</i> (<i>pine</i>)	Ethanollic and aqueous extracts	STZ-induced diabetic rats/ (GSH and MDA in liver, kidney, and heart tissues)	[98]
		<i>V. album</i> / <i>Acer campestre</i> ; <i>Fraxinus excelsio</i> ; <i>Populus nigra</i> ; <i>Malus domestica</i> ; <i>Robinia pseudoacacia</i>	Aqueous extract	In vitro (DPPH, ORAC, and TEAC assay)	[65]
		<i>V. coloratum</i> /ns*	Ethanollic extract	In vitro (Hydroxyl Radical Scavenging Assay; Superoxide Anion Radical Scavenging Assay)	[99]
	Antiepileptic	<i>V. album</i> L./ Citrus	Aqueous extract	Swiss albino mice and Wistar albino rats (MES-induced seizure; INH-induced convulsions)	[100]

*ns- not specified; MDA- malondialdehyde; MES- Maximum electroshock; INH-Isoniazid;

Utilization of mistletoe extract in medicine, cosmetics, and functional products

Patents issued from 2011 to 2021 related to mistletoe extract (*V. album*) and its applications in medical treatments, cosmetic formulations, and functional products are outlined in **Table 4**.

Table 4. List of patents published from on *V. album* (2011-2021)

Patent no.	Publication date	Title	Purpose
KR20110136539A	2011-12-21	Korean mistletoe extracts have antiobesity and hepatic steatosis protection activity	- Reduces obesity by preventing fat accumulation in the epididymal region and liver.
KR20140115730A	2014-10-31	Shampoo composition for preventing hair loss and promoting hair growth comprising extracts of <i>V. album</i> and <i>Chamaecyparis obtusa</i>	-Helps combat hair loss. - Demonstrates enhanced antioxidant properties.
CN104707097A	2015-06-17	Pharmaceutical composition <i>Viscum</i> and astragali radix powder and use thereof in the preparation of medicines for blocking precancerous lesions of the liver and treating liver cancer or viral hepatitis	- Lowers the incidence of GGT foci in early-stage precancerous lesions. - Exhibits hydroxyl radical scavenging activity, preventing the initiation and progression of liver precancerous lesions.
KR20160017903A	2016-02-17	<i>V. album</i> extract containing a sweet jelly preparation method and sweet jelly prepared by the method	- Strengthens the immune system. - Suggested for use in immunotherapy and cancer prevention.
KR20170053544A	2017-05-16	Cosmetic composition for moisturizing and anti-aging containing extracts of <i>Viscum album</i> , <i>Chamaecyparis obtusa</i> , <i>Quercus Robur</i> , and <i>Camellia japonica</i> Linne	- Acts as a skin astringent. - Minimizes cytotoxic effects triggered by environmental stressors. - Regulates inflammatory mediator expression.
CN106692909A	2017-05-24	<i>Pyrus pyrifolia</i> tree <i>Viscum coloratum</i> grease with lung-clearing and phlegm-eliminating effects	- Aids in lung detoxification and phlegm clearance.

CN108295100A	2018-07-20	Application of the <i>V. album</i> extract in preparing drugs for rheumatoid arthritis	- Inhibits inflammatory mediator synthesis by targeting NF-κB signaling pathways. - Reduces inflammation in the synovial membrane. - Protects against joint and bone degradation.
KR102007096B1	2019-08-02	Composition for preventing or treating hearing loss comprising <i>Viscum ovalifolium</i> extracts	- Helps prevent hearing impairment.
KR20200109068A	2020-09-22	A soap comprising <i>V. album</i> fermentation product and method for preparing the same	- Rapidly penetrates the skin. - Soothes skin inflammation. - Provides an exfoliating effect, eliminating harmful substances from the skin's surface. - Delays skin aging due to the antioxidant components in <i>V. album</i>.
BR102020003563A2	2021-09-08	Topical pharmaceutical composition with antitumor activity containing <i>V. album</i>	- Exhibits antitumor properties.

GGT gamma-glutamyltranspeptidase, GPT glutamic—pyruvic transaminase, NF- κ B Nuclear factor- κ B

Conclusion

Mistletoe demonstrates significant therapeutic potential for various health conditions. Its unique phytochemical makeup and corresponding biological effects are influenced by the host tree from which it grows. Among its key bioactive compounds, lectins and viscotoxins play a crucial role in its anticancer properties. Additionally, mistletoe contains an array of beneficial secondary metabolites, including flavonoids and phenolic acids, which are widely distributed in plants. There is strong evidence suggesting that the consumption of these compounds can contribute to a reduction in the incidence of cancer, cardiovascular diseases, and diabetes. Further research into the interactions between mistletoe polyphenols and lectins is essential to uncover new therapeutic applications of mistletoe-based treatments.

Acknowledgments: We would like to acknowledge the support of the University of Oradea for funding this project through the “Excellent Scientific Research Related to Priority Fields with the Goal of Technology Transfer: INO-TRANSFER-UO” competition, Project number 309/21.12.2021.

Conflict of Interest: None

Financial Support: This research was supported by the European Union under the Horizon 2020 “NextFood” project, Grant Agreement No. 771738.

Ethics Statement: None

References

1. Amico GC, Nickrent DL, Vidal-Russell R. Macroscale analysis of mistletoe host ranges in the Andean-Patagonian forest. *Plant Biol Stuttg Ger.* 2019;21(1):150-6.
2. Szurpnicka A, Zjawiony JK, Szterk A. Therapeutic potential of mistletoe in CNS-related neurological disorders and the chemical composition of *Viscum* species. *J Ethnopharmacol.* 2019;231:241-52.
3. Combes C. Parasitism: the ecology and evolution of intimate interactions (interspecific interactions). *Q Rev Biol.* The University of Chicago Press; 2002;77:207. doi:10.1086/343967
4. Krasnylenko Y, Těšitel J, Ceccantini G, Oliveira-da-Silva M, Dvořák V, Steele D, et al. Parasites on parasites: hyper-, epi, and autoparasitism among flowering plants. *Am J Bot.* 2021;108(1):8-21.
5. Böhling N, Greuter W, Raus T, Snogerup B, Snogerup S, Snogerup B. Notes on the Cretan mistletoe, *Viscum album* subsp. *creticum* subsp. *nova* (Loranthaceae/Viscaceae). *Isr J Plant Sci.* 2002;50(sup1):77-84.

6. Costa JPL, Brito HO, Galvão-Moreira LV, Brito LGO, Costa-Paiva L, Brito LMO. Randomized double-blind placebo-controlled trial of the effect of *Morus nigra* L. (black mulberry) leaf powder on symptoms and quality of life among climacteric women. *Int J Gynaecol Obstet.* 2020;148(2):243-52.
7. Zuber D. Biological flora of central europe: *Viscum album* L. *Flora - Morphol Distrib Funct Ecol Plants.* 2004;199(3):181-203. Available from: <https://www.sciencedirect.com/science/article/pii/S0367253004701103>
8. Peñaloza E, Holandino C, Scherr C, Araujo PIP de, Borges RM, Urech K, et al. Comprehensive metabolome analysis of fermented aqueous extracts of *Viscum album* L. by liquid chromatography-high resolution tandem mass spectrometry. *Mol Basel Switz.* 2020;25(17):E4006.
9. Vergara-Barberán M, Lerma-García MJ, Nicoletti M, Simó-Alfonso EF, Herrero-Martínez JM, Fasoli E, et al. Proteomic fingerprinting of mistletoe (*Viscum album* L.) via combinatorial peptide ligand libraries and mass spectrometry analysis. *J Proteomics.* 2017;164:52-8.
10. Bonamin LV, De Carvalho AC, Waisse S. *Viscum album* (L.) in experimental animal tumors: a meta-analysis. *Exp Ther Med.* 2017;13(6):2723-40. Available from: <https://www.spandidos-publications.com/10.3892/etm.2017.4372>
11. Kwon YS, Chun SY, Kim MK, Nan HY, Lee C, Kim S. Mistletoe extract targets the STAT3-FOXO1 pathway to induce apoptosis and inhibits metastasis in breast cancer cells. *Am J Chin Med.* 2021;49(02):487-504. doi:10.1142/S0192415X21500221
12. Pietrzak W, Nowak R, Gawlik-Dziki U, Lemieszek M, Rzeski W. LC-ESI-MS/MS identification of biologically active phenolic compounds in mistletoe berry extracts from different host trees. *Molecules.* 2017;22(4):624.
13. Pietrzak W, Nowak R. Impact of harvest conditions and host tree species on chemical composition and antioxidant activity of extracts from *Viscum album* L. *Mol Basel Switz.* 2021;26(12):3741.
14. Fritz P, Dippon J, Müller S, Goletz S, Trautmann C, Pappas X, et al. Is Mistletoe treatment beneficial in invasive breast cancer? A new approach to an unresolved problem. *Anticancer Res.* 2018;38(3):1585-93.
15. Rostock M. Mistletoe in the treatment of cancer patients. *Bundesgesundheitsblatt Gesundheitsforschung Gesundheitsschutz.* 2020;63(5):535-40.
16. Wode K, Hök Nordberg J, Kienle GS, Elander NO, Bernhardson BM, Sunde B, et al. Efficacy of mistletoe extract as a complement to standard treatment in advanced pancreatic cancer: study protocol for a multicentre, parallel-group, double-blind, randomized, placebo-controlled clinical trial (MISTRAL). *Trials.* 2020;21(1):783.
17. Shakeel M, Trinidad A, Geider S, Ah-See KW. The case for mistletoe in the treatment of laryngeal cancer. *J Laryngol Otol.* 2014;128(3):302-6.
18. Rexer H, Geschäftsstelle der AUO. Study for the treatment of nonmuscle invasive bladder cancer: a phase III efficacy trial for intravesical instillation of mistletoe extract in superficial bladder cancer (TIM) - AB 40/11 of the AUO. *Urol Ausg A.* 2016;55(7):963-5.
19. Park YK, Do YR, Jang BC. Apoptosis of K562 leukemia cells by Abnobaviscum F®, a European mistletoe extract. *Oncol Rep.* 2012;28(6):2227-32. Available from: <https://www.spandidos-publications.com/10.3892/or.2012.2026>
20. Menke K, Schwermer M, Eisenbraun J, Schramm A, Zuzak TJ. Anticancer effects of *Viscum album* fraxini extract on medulloblastoma cells in vitro. *Complement Med Res.* 2021;28(1):15-22.
21. Schötterl S, Miemietz JT, Ilina EI, Wirsik NM, Ehrlich I, Gall A, et al. Mistletoe-based drugs work in synergy with radio-chemotherapy in the treatment of glioma in vitro and in vivo in glioblastoma bearing mice. *Evid-Based Complement Altern Med ECAM.* 2019;2019:1376140.
22. Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et al. The PRISMA 2020 statement: an updated guideline for reporting systematic reviews. *BMJ.* 2021;372:n71. Available from: <https://www.bmj.com/content/372/bmj.n71>
23. Halbritter DA, Willett DS, Gordon JM, Stelinski LL, Daniels JC. Behavioral evidence for host transitions in plant, plant parasite, and insect interactions. *Environ Entomol.* 2018;47(3):646-53.
24. Liu B, Bian H, Bao J. Plant lectins: Potential antineoplastic drugs from bench to clinic. *Cancer Lett.* 2010;287(1):1-12. Available from: <https://www.sciencedirect.com/science/article/pii/S0304383509003450>
25. Liu B, Le CT, Barrett RL, Nickrent DL, Chen Z, Lu L, et al. Historical biogeography of Lorantheae (Santalales): diversification agrees with the emergence of tropical forests and radiation of songbirds. *Mol*

- Phylogenet Evol. 2018;124:199-212. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S1055790317308217>
26. Baena-Díaz F, Ramírez-Barahona S, Ornelas JF. Hybridization and differential introgression are associated with environmental shifts in a mistletoe species complex. *Sci Rep.* 2018;8(1):5591.
27. Maclean AE, Hertle AP, Ligas J, Bock R, Balk J, Meyer EH. Absence of complex i is associated with diminished respiratory chain function in European mistletoe. *Curr Biol.* 2018;28(10):1614-9. Available from: <https://linkinghub.elsevier.com/retrieve/pii/S0960982218303658>
28. Fu J, Wan L, Song L, He L, Jiang N, Long H, et al. Identification of MicroRNAs in *Taxillus Chonensis* (DC.) Danser Seeds under Cold Stress. *BioMed Res Int.* 2021;2021:5585884.
29. Oei SL, Thronicke A, Schad F. Mistletoe and immunomodulation: insights and implications for anticancer therapies. *Evid-Based Complement Altern Med ECAM.* 2019;2019:5893017.
30. Jones P. The Tenbury mistletoe festival & national mistletoe day. *Holidappy.* [cited 2022 Jan 16]. Available from: <https://holidappy.com/holidays/The-Tenbury-Mistletoe-Festival>
31. Ostermann T, Appelbaum S, Poier D, Boehm K, Raak C, Büssing A. A systematic review and meta-analysis on the survival of cancer patients treated with a fermented *Viscum album* L. extract (Iscador): an update of findings. *Complement Med Res.* 2020;27(4):260-71.
32. Ostermann T, Büssing A. Retrolective studies on the survival of cancer patients treated with mistletoe extracts: a meta-analysis. *Explore N Y N.* 2012;8(5):277-81.
33. Jäger T, Holandino C, Melo MN de O, Peñaloza EMC, Oliveira AP, Garrett R, et al. Metabolomics by UHPLC-Q-TOF reveals host tree-dependent phytochemical variation in *Viscum album* L. plants. 2021;10(8):1726. Available from: <https://www.mdpi.com/2223-7747/10/8/1726>
34. Ramm H, Urech K, Scheibler M, Grazi G. Cultivation and development of *Viscum album* L. Mistletoe. CRC Press; 2000. p. 91-110.
35. Maul K, Krug M, Nickrent DL, Müller KF, Quandt D, Wicke S. Morphology, geographic distribution, and host preferences are poor predictors of phylogenetic relatedness in the mistletoe genus *Viscum* L. *Mol Phylogenet Evol.* 2019;131:106-15.
36. Büssing A. Mistletoe: the Genus *Viscum*. CRC Press; 2000.
37. Wilson CA, Calvin CL. An origin of aerial branch parasitism in the mistletoe family, Loranaceae. *Am J Bot.* 2006;93(5):787-96. doi:10.3732/ajb.93.5.787
38. Barney C, Hawksworth F, Geils B. Hosts of *Viscum album*. *Eur J for Pathol.* 2007;28(3):187-208.
39. Danser DG. Long. *Viscum album* subsp. *meridian*. *Plants World Online.* 1982. Available from: <http://powo.science.kew.org/taxon/urn:lsid:ipni.org:names:921668-1>
40. Long C, Fan R, Zhang Q, Zhang Z, Wang D, Xia Y, et al. Simultaneous identification and quantification of the common compounds of *Viscum coloratum* and its corresponding host plants by ultra-high performance liquid chromatography with quadrupole time-of-flight tandem mass spectrometry and triple quadrupole mass spectrometry. *J Chromatogr B Analyt Technol Biomed Life Sci.* 2017;1061:176-84. doi:10.1016/j.jchromb.2017.07.028
41. Lyu SY, Park SM, Choung BY, Park WB. Comparative study of Korean (*Viscum album* var. *coloratum*) and European mistletoes (*Viscum album*). *Arch Pharm Res.* 2000;23(6):592-8. doi:10.1007/BF02975247
42. Patel BP, Singh PK. *Viscum articulatum* Burm. f.: a review on its phytochemistry, pharmacology and traditional uses. *J Pharm Pharmacol.* 2018;70(2):159-77. doi:10.1111/jphp.12837
43. Kunwar R, Adhikari N, Devkota M. Indigenous use of mistletoes in tropical and temperate region of Nepal. *Banko Janakari.* 2007;15(2):38-42.
44. Deeni Y, Sadiq N. Antimicrobial properties and phytochemical constituents of the leaves of African mistletoe (*Tapinanthus dodoneifolius* (DC) Danser) (Loranthaceae): an ethnomedicinal plant of Hausaland, Northern Nigeria. *J Ethnopharmacol.* 2003;83(3):235-40.
45. Obatomi D, Bikomo EO, Temple V. Anti-diabetic properties of the African mistletoe in streptozotocin-induced diabetic rats. *J Ethnopharmacol.* 1994;43(1):13-7.
46. Clark NF, McComb JA, Taylor-Robinson AW, Clark NF, McComb JA, Taylor-Robinson AW. Host species of mistletoes (Loranthaceae and Viscaceae) in Australia. *Aust J Bot.* 2020;68(1):1-13.
47. Govaerts R, Nic Lughadha E, Black N, Turner R, Paton A. The world checklist of vascular plants, a continuously updated resource for exploring global plant diversity. *Sci Data.* 2021;8(1):215. Available from: <https://www.nature.com/articles/s41597-021-00997-6>

48. Watson DM, Herring M. Mistletoe as a keystone resource: an experimental test. *Proc Biol Sci.* 2012;279(1743):3853-60.
49. Watson DM. Mistletoe—A keystone resource in forests and woodlands worldwide. *Annu Rev Ecol Syst.* 2001;32(1):219-49. doi:10.1146/annurev.ecolsys.32.081501.114024
50. Cordero CM, Gil Serrano AM, Ayuso Gonzalez MJ. Transfer of bipiperidyl and quinolizidine alkaloids to *Viscum cruciatum* Sieber (Loranthaceae) hemiparasitic on *Retama sphaerocarpa* boissier (Leguminosae). *J Chem Ecol.* 1993;19(10):2389-93. doi:10.1007/BF00979672
51. Zhang RZ, Zhao JT, Wang WQ, Fan RH, Rong R, Yu ZG, et al. Metabolomics-based comparative analysis of the effects of host and environment on *Viscum coloratum* metabolites and antioxidative activities. *J Pharm Anal.* 2022;12(2):243-52. Available from: <https://www.sciencedirect.com/science/article/pii/S2095177921000472>
52. Kaurinovic B, Vastag D. Flavonoids and phenolic acids as potential natural antioxidants. IntechOpen; 2019. Available from: <https://www.intechopen.com/chapters/65331>
53. Fritea L, Ganea M, Zdrinca M, Dobjanschi L, Antonescu A, Vicas SI, et al. Perspectives on the combined effects of *Ocimum basilicum* and *Trifolium pratense* extracts in terms of phytochemical profile and pharmacological effects. *Plants.* 2021;10(7):1390.
54. Vicas SI, Cavalu S, Laslo V, Tocai M, Costea TO, Moldovan L. Growth, photosynthetic pigments, phenolic, glucosinolates content and antioxidant capacity of broccoli sprouts in response to nanoselenium particles supply. *Not Bot Horti Agrobot Cluj-Napoca.* 2019;47(3):821-8.
55. Alabdallat NG. In-vivo antioxidant effects of the orally administered paracetamol, aqueous extracts of *Salvia triloba*, and *Origanum syriacum*. *J Biochem Technol.* 2021;12(4):19-22. Available from: <https://jbiochemtech.com/article/in-vivo-antioxidant-effects-of-the-orally-administered-paracetamol-aqueous-extracts-of-salvia-trilo-g3huxydyndagsa>
56. Al-Harbi D, Awad N, Alsberi H, Abdein M. Apoptosis induction, cell cycle arrest and in vitro anticancer potentiality of *Convolvulus spicatus* and *Astragalus vogelii*. *World J Environ Biosci.* 2020;8(4):69-75. Available from: www.environmentaljournal.org
57. Darwati D, Safitri AN, Ambardhani N, Mayanti T, Nurlelarsi N, Kurnia D. Effectiveness and anticancer activity of a novel phenolic compound from *Garcinia porrecta* against the MCF-7 breast cancer cell line in vitro and in silico. *Drug Des Devel Ther.* 2021;15:3523-33. Available from: <https://www.dovepress.com/effectiveness-and-anticancer-activity-of-a-novel-phenolic-compound-fro-peer-reviewed-fulltext-article-DDDT>
58. Alotaibi BS, Ijaz M, Buabeid M, Kharaba ZJ, Yaseen HS, Murtaza G. Therapeutic effects and safe uses of plant-derived polyphenolic compounds in cardiovascular diseases: a review. *Drug Des Devel Ther.* 2021;15:4713-32. Available from: <https://www.dovepress.com/therapeutic-effects-and-safe-uses-of-plant-derived-polyphenolic-compou-peer-reviewed-fulltext-article-DDDT>
59. Vicas S, Prokisch J, Rugina OD, Socaciu C. Hydrophilic and lipophilic antioxidant activities of mistletoe (*Viscum album*) as determined by FRAP method. *Not Bot Horti Agrobot Cluj-Napoca.* 2009;37(2):112-6.
60. Vicas SI, Rugin D, Socaciu C. Comparative study about antioxidant activities of *Viscum album* from different host trees, harvested in different seasons. *J Med Plants Res.* 2011;5(11):2237-44. Available from: <https://academicjournals.org/journal/JMPR/article-abstract/4B70C8121233>
61. Olanlokun JO, Ekundayo MT, Ebenezer O, Koorbanally NA, Olorunsogo OO. Antimalarial and erythrocyte membrane stability properties of *Globimetula braunii* (engle van Tiegh) growing on cocoa in plasmodium berghei-infected mice. *Infect Drug Resist.* 2021;14:3795-808. Available from: <https://www.dovepress.com/antimalarial-and-erythrocyte-membrane-stability-properties-of-globimet-peer-reviewed-fulltext-article-IDR>
62. Kumar S, Pandey AK. Chemistry and biological activities of flavonoids: an overview. *Sci World J.* 2013;2013:162750.
63. Vicas SI, Rugin D, Socaciu C. Comparative study about antioxidant activities of *Viscum album* from different host trees, harvested in different seasons. *J Med Plants Res.* 2011;5(11):2237-44.
64. Luczkiewicz M, Cisowski W, Kaiser P, Ochocka R, Piotrowski A. Comparative analysis of phenolic acids in mistletoe plants from various hosts. *Acta Pol Pharm.* 2001;58(5):373-9.

65. Vicaș SI, Rugina OD, Leopold L, Pinte A, Socaciu C. HPLC fingerprint of bioactive compounds and antioxidant activities of *Viscum album* from different host trees. *Not Bot Horti Agrobot Cluj-Napoca*. 2011;39(1):48-57. Available from: <https://www.notulaeobotanicae.ro/index.php/nbha/article/view/3455>
66. Haas K, Bauer M, Wollenweber E. Cuticular waxes and flavonol aglycones of mistletoes. *Z Für Naturforschung C J Biosci*. 2003;58(7-8):464-70. Available from: <https://eurekamag.com/research/003/699/003699122.php>
67. Stefanucci A, Zengin G, Llorent-Martínez E, Dimmito M, Della Valle A, Pieretti S, et al. *Viscum album* L. homogenizer-assisted and ultrasound-assisted extracts as potential sources of bioactive compounds. *J Food Biochem*. 2020;44(9):e13377.
68. Urech K, Baumgartner S. Chemical constituents of *Viscum album* L.: Implications for the pharmaceutical preparation of mistletoe. *Mistletoe: From mythology to evidence-based medicine*. 2015;4:11-23.
69. Elluru SR, Duong Van Huyen JP, Delignat S, Prost F, Bayry J, Kazatchkine M, et al. Molecular mechanisms underlying the immunomodulatory effects of mistletoe (*Viscum album* L.) extracts *Iscador*: Review. *Arzneimittelforschung*. 2006;56(06):461-6.
70. Peumans WJ, Van Damme JM, Barre A, Rougé P. Classification of plant lectins in families of structurally and evolutionary related proteins. In: Wu AM, editor. *Molecular Immunology of Complex Carbohydrates — 2*. Boston (MA): Springer US; 2001. p. 27–54. doi:10.1007/978-1-4615-1267-7_3
71. Peumans W, Verhaert P, Pfüller U, Damme EV. Isolation and partial characterization of a small chitin-binding lectin from mistletoe (*Viscum album*). *FEBS Lett*. 1996;396(2-3):261-5.
72. Wacker R, Stoeva S, Pfüller K, Pfüller U, Voelter W. Complete structure determination of the A chain of mistletoe lectin III from *Viscum album* L. ssp. *album*. *J Pept Sci Off Publ Eur Pept Soc*. 2004;10(3):138-48.
73. Mazalovska M, Kouokam JC. Transiently expressed mistletoe lectin II in *Nicotiana benthamiana* demonstrates anticancer activity in vitro. *Mol Basel Switz*. 2020;25(11):E2562.
74. Kang TB, Song SK, Yoon TJ, Yoo YC, Lee KH, Her E, et al. Isolation and characterization of two Korean mistletoe lectins. *BMB Rep*. 2007;40(6):959-65. Available from: <http://www.koreascience.or.kr/article/JAKO200710103429562.page>
75. Endo Y, Tsurugi K, Franz H. The site of action of the A-chain of mistletoe lectin I on eukaryotic ribosomes. The RNA N-glycosidase activity of the protein. *FEBS Lett*. 1988;231(2):378-80.
76. Soler MH, Stoeva S, Schwamborn C, Wilhelm S, Stiefel T, Voelter W. Complete amino acid sequence of the a chain of mistletoe lectin I. *FEBS Lett*. 1996;399(1-2):153-7. doi:10.1016/s0014-5793(96)01309-9
77. Wacker R, Stoeva S, Betzel C, Voelter W. Complete structure determination of N-acetyl-D-galactosamine-binding mistletoe lectin-3 from *Viscum album* L. *album*. *J Pept Sci Off Publ Eur Pept Soc*. 2005;11(6):289-302.
78. Beztsinna N, de Matos MBC, Walther J, Heyder C, Hildebrandt E, Lenewit G, et al. Quantitative analysis of receptor-mediated uptake and pro-apoptotic activity of mistletoe lectin-1 by high content imaging. *Sci Rep*. 2018;8(1):2768.
79. Franz H. Mistletoe lectins and their A and B chains. *Oncology*. 1986;43(Suppl 1):23-34.
80. Bohlmann H, Apel K. Thionins. *Annu Rev Plant Biol*. 1991;42(1):227-40.
81. Schaller G, Urech K, Grazi G, Giannattasio M. Viscotoxin composition of the three European subspecies of *Viscum album*. *Planta Med*. 1998;64(07):677-8. doi:10.1055/s-2006-957553
82. Cavalu S, Damian G. Rotational correlation times of 3-carbamoyl-2, 2, 5, 5-tetramethyl-3-pyrrolin-1-yloxy spin label with respect to heme and nonheme proteins. *Biomacromolecules*. 2003;4(6):1630-5.
83. Karagöz A, Önay E, Arda N, Kuru A. Antiviral potency of mistletoe (*Viscum album* ssp. *album*) extracts against human parainfluenza virus type 2 in Vero cells. *Phytother Res*. 2003;17(5):560-2.
84. Kleinsimon S, Kauczor G, Jaeger S, Eggert A, Seifert G, Delebinski C. *Viscum*TT induces apoptosis and alters IAP expression in osteosarcoma in vitro and has synergistic action when combined with different chemotherapeutic drugs. *BMC Complement Altern Med*. 2017;17(1):1-3.
85. Han SY, Hong CE, Kim HG, Lyu SY. Anti-cancer effects of enteric-coated polymers containing mistletoe lectin in murine melanoma cells in vitro and in vivo. *Mol Cell Biochem*. 2015;408(1):73-87.
86. Tröger W, Galun D, Reif M, Schumann A, Stanković N, Milićević M. *Viscum album* [L.] extract therapy in patients with locally advanced or metastatic pancreatic cancer: a randomized clinical trial on overall survival. *Eur J Cancer*. 2013;49(18):3788-97. Available from: [https://www.cabdirect.org/cabdirect/abstract/20143031671?q=\(au%3A%22Reif%2C+M.%22\)](https://www.cabdirect.org/cabdirect/abstract/20143031671?q=(au%3A%22Reif%2C+M.%22))

87. Werthmann P, Sträter G, Friesland H, Kienle G. Durable response of cutaneous squamous cell carcinoma following high-dose peri-lesional injections of *Viscum album* extracts – a case report. *Phytomedicine Int J Phytother Phytopharm.* 2013;20(3-4):324-7.
88. Ofem OE, Eno AE, Imoru J, Nkanu E, Unoh F, Ibu JO. Effect of crude aqueous leaf extract of *Viscum album* (mistletoe) in hypertensive rats. *Indian J Pharmacol.* 2007;39(1):15. Available from: <https://www.ijp-online.com/article.asp?issn=0253-7613;year=2007;volume=39;issue=1;spage=15;epage=19;aulast=Ofem;type=0>
89. Radenkovic M, Ivetic V, Popovic M, Brankovic S, Gvozdenovic L. Effects of mistletoe (*Viscum album* L., Loranaceae) extracts on arterial blood pressure in rats treated with atropine sulfate and hexocycline. *Clin Exp Hypertens.* 2009;31(1):11-9. Available from: [https://www.cabdirect.org/cabdirect/abstract/20093082399?q=\(au%3a%22Radenkovic%2c+M.%22\)](https://www.cabdirect.org/cabdirect/abstract/20093082399?q=(au%3a%22Radenkovic%2c+M.%22))
90. Poruthukaren KJ, Palatty PL, Baliga MS, Suresh S. Clinical evaluation of *Viscum album* mother tincture as an antihypertensive: a pilot study. *J Evid-Based Complement Altern Med.* 2014;19(1):31-5.
91. Bachhav SS, Patil SD, Bhutada MS, Surana SJ. Oleanolic acid prevents glucocorticoid-induced hypertension in rats. *Phytother Res.* 2011;25(10):1435-9. doi:10.1002/ptr.3431
92. Shahabuddin M, Pouramir M, Moghadamnia A, Lakzaei M, Mirhashemi SM, Motallebi M. Antihyperglycemic and antioxidant activity of *Viscum album* extract. *Afr J Pharm Pharmacol.* 2011;5(3):433-6.
93. Adaramoye O, Amanlou M, Habibi-Rezaei M, Pasalar P, Ali MM. Methanolic extract of African mistletoe (*Viscum album*) improves carbohydrate metabolism and hyperlipidemia in streptozotocin-induced diabetic rats. *Asian Pac J Trop Med.* 2012;5(6):427-33.
94. Ertürk Ö, Kati H, Yayli N, Demirbag Z. Antimicrobial activity of *Viscum album* L. subsp. *abietis* (Viesb). *Turk J Biol.* 2003;27(4):255-8. Available from: <https://app.trdizin.gov.tr/makale/TXpBME1qYzM=/antimicrobial-activity-of-viscum-album-l-subsp-abietis-viesb->
95. Orhue PO, Edomwande EC, Ighinosa E, Momoh ARM, Asekomhe OO. Antibacterial activity of extracts of mistletoe (*Tapinanthus dodoneifolius* (Dc) Dancer) from cocoa tree (*Theobroma cacao*). *Int J Herbs Pharmacol Res.* 2014;3(1):24-9. Available from: <https://www.ajol.info/index.php/ijhpr/article/view/104727>
96. Yusuf L, Oladunmoye MK, Ogundare AO, Mohmo MO, Daudu OA. Comparative antifungal and toxicological effects of the extract of mistletoes growing on two different host plants in Akure North, Nigeria. *Int J Biotechnol Food Sci.* 2014;2(2):31-4. Available from: <http://sciencewebpublishing.net/ijbfs/archive/2014/Feb/Abstract/Yusuf%20et%20al.htm>
97. Deliorman Orhan D. Evaluation of antimycobacterial activity of *Viscum album* subspecies. *Pharm Biol.* 2001;39(5):381-3.
98. Orhan DD, Aslan M, Sendogdu N, Ergun F, Yesilada E. Evaluation of the hypoglycemic effect and antioxidant activity of three *Viscum album* subspecies (European mistletoe) in streptozotocin-diabetic rats. *J Ethnopharmacol.* 2005;98(1-2):95-102.
99. Yao H, Liao ZX, Wu Q, Lei GQ, Liu ZJ, Chen DF, et al. Antioxidative flavanone glycosides from the branches and leaves of *Viscum coloratum*. *Chem Pharm Bull (Tokyo).* 2006;54(1):133-5. doi:10.1248/cpb.54.133
100. Gupta G, Kazmi I, Afzal M, Rahman M, Saleem S, Ashraf MS, et al. Sedative, antiepileptic and antipsychotic effects of *Viscum album* L. (Loranthaceae) in mice and rats. *J Ethnopharmacol.* 2012;141(3):810-6. Available from: <https://www.sciencedirect.com/science/article/pii/S0378874112001717>